Fax: 832-895-4040
Phone: 832-895-5000
E-mail: intake@adaptivinfusion.com

adaptIV infusion

Leqembi™ (Lecanemab-irmb)

Provider Order Form rev. 8/10/2023

PATIENT INFORMATION

Referral Status *(check one):* ☐ New Referral ☐ Updated Order ☐ Order Renewal

Patient Name: _____ DOB: _____

NKDA ☐ Allergies: _____ Weight _____ *Please specify:* ☐ lbs ☐ kg Height: _____

Patient Status *(check one):* ☐ New to Therapy ☐ Continuing Therapy | Last Treatment Date: _____ Next Due Date: _____

ICD-10 code (required): _____ ICD-10 description: _____

REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.

PRESCRIPTION

☐ Provide nursing care per AdaptIV Infusion Nursing Procedures, including reaction management and post-procedure observation
☐ Medicare Registry # _____

DIAGNOSIS:
☐ G30.0 Alzheimer's Disease, Early Onset
☐ G30.1 Alzheimer's Disease, Late Onset
☐ G30.8 Other Alzheimer's disease
☐ G30.9 Alzheimer's disease, unspecified
☐ G31.84 Mild Cognitive Impairment, So Stated

G30.X codes require secondary F02.8X code BELOW:
☐ F02.80 Dementia without behavioral disturbance
☐ F02.81 Dementia with behavioral disturbance

PRESCRIBER MUST INDICATE THE FOLLOWING REQUIREMENTS HAVE BEEN MET (PLEASE PROVIDE DOCUMENTATION):
☐ Beta Amyloid Pathology Confirmed Via:
☐ Amyloid PET Scan Date: _____
OR
☐ CSF Analysis Date: _____ Result: _____
☐ Cognitive Assessment Used: _____
Date: _____ Result: _____
☐ ApoE E4 Genetic Test: Date: _____
Result: ☐ Homozygote ☐ Heterozygote ☐ Noncarrier

PRE-INFUSION:
☐ Confirm baseline MRI results prior to initiation of treatment.
☐ Confirm MRI completed and reviewed by prescriber prior to the 5th, 7th, and 14th treatment.
☐ Measure and record weight prior to each treatment to determine dose.
☐ Hold infusion and notify provider if patient reports:
- Headache
- Vision changes
- Dizziness
- New or worsening confusion
- Nausea

MEDICATION:
☐ Administer LEQEMBI 10 mg/kg intravenously over at least 60 minutes.
☐ Dilute required volume of lecanemab-irmb in 250 ml 0.9% sodium chloride and infuse using a terminal low-protein binding 0.2-micron in-line filter.
☐ If infusion-related reaction occurs, stop infusion and treat per orders/protocol as clinically indicated.

TREATMENT FREQUENCY:
☐ 10mg/kg IV every 2 weeks
☐ 10mg/kg IV every 4 weeks (after 18 months of treatment only)

POST-INFUSION:
☐ Educate patient/care partner to report headache, dizziness, nausea, vision changes, or new/worsening confusion.
☐ Fax treatment notes to provider at number below.

SPECIAL INSTRUCTIONS

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip Code: _____

_____ _____ _____
Provider Name (Print) *Provider Signature* *Date*

REQUIRED: PLEASE INCLUDE ALL REQUIRED LABS AND A COPY OF PATIENT'S INSURANCE CARD – FRONT AND BACK